Pelvic Floor Pre-Procedure Orders

To be performed within 30 days, unless otherwise noted.
Fax to (513) 585-0169

Surgeon Name: _____
Phone: _____
Fax: _____

Patient Name : _____

Date of Birth : _____

Surgery confirmation # _____

Procedure Orders: _____

_____

WEIGHT (kg): _____ ALLERGIES: _____

Before Day of Procedure:

Labs:

- ☐ CBC
- ☐ HGBA1C
- ☐ PT/INR
- ☐ PTT
- ☐ Basic Metabolic Panel (EP1)

Day of Procedure:

Nursing Orders:

- ☒ Obtain Vital Signs Pre and Post Procedure
- ☒ POC Urine Pregnancy Test (For Fluoroscopy cases), if Female with no history of hysterectomy and less than 55 years of age
- ☐ Nitrous Oxide inhalation self-administered via Nitrouseal system. Start at 30% Nitrous Oxide and 70% oxygen blend. Increase by 5% every 1-2 minutes until desired patient analgesia is achieved. The maximum concentration is 50% Nitrous Oxide.

Medications:

- ☐ No Pre op Antibiotic Necessary

Pelvic floor disorder Antibiotics:

- ☐ CephALEXin (Keflex) capsule, 500 mg, oral X1
- ☐ For PCN or Cephalosporin allergy: levoFLOXacin (Levaquin) tablet, 750 mg, oral X1

Prostate Biopsy Antibiotics:

- ☐ CephALEXin (Keflex) capsule, 500 mg, oral X1
- ☐ For PCN or Cephalosporin allergy: levoFLOXacin (Levaquin) tablet, 750 mg, oral X1
- ☐ Gentamicin Intramuscular, 80 mg intramuscular X1 pre-procedure

Botox and other Medications:

- ☐ Botulinum toxin type A (BOTOX) injection for Bladder, 100 units X1
- ☐ Botulinum toxin type A (BOTOX) injection for Bladder, 200 units X1
- ☐ Botulinum toxin type A (BOTOX) injection for Rectal, 50 units X1
- ☐ Phenazopyridine (PYRIDIUM) tablet oral, 200 mg X1
- ☐ Triamcinolone (KENALOG) 40 mg/ml - Dilute with 9 ml of sodium chloride, 0.9% in a 10 ml syringe. Will be administered cystoscopically via injection needle.

OTHER ORDERS:

Physician Signature _____ Date: _____ Time: _____

R3854